

CONSTRUCTION OF EXOTIC SMOOTH STRUCTURES

ANAR AKHMEDOV

ABSTRACT. In this article, we construct infinitely many simply connected, nonsymplectic and pairwise nondiffeomorphic 4-manifolds starting from the elliptic surfaces $E(n)$ and applying the sequence of knot surgery, ordinary blowups and rational blowdown. We also compute the Seiberg-Witten invariants of these manifolds.

0. INTRODUCTION

There has been a recent flurry of activity in the discovery of exotic smooth structures on simply-connected 4-manifolds with small Euler characteristic. In the 2004, Jongil Park [P2] gave the first example of exotic smooth structure on $\mathbf{CP}^2 \# 7\overline{\mathbf{CP}}^2$ i.e. 4-manifold homeomorphic to $\mathbf{CP}^2 \# 7\mathbf{CP}^2$ but not diffeomorphic to it. Shortly afterwards, András Stipsicz and Zoltán Szabó used a technique similar to Park's to construct an exotic smooth structure on $\mathbf{CP}^2 \# 6\overline{\mathbf{CP}}^2$ [SS1]. Then Fintushel and Stern [FS3] introduced a new technique, surgery in double nodes, which demonstrated that in fact $\mathbf{CP}^2 \# k\overline{\mathbf{CP}}^2$, $k = 6, 7, 8$, have infinitely many distinct smooth structures. Park, Stipsicz, and Szabó [PSS], using [FS3], constructed infinitely many smooth structures when $k = 5$. Stipsicz and Szabó used similar ideas to construct exotic smooth structures on $3\mathbf{CP}^2 \# k\overline{\mathbf{CP}}^2$ for $k = 9$ [SS2] and Park for $k = 8$ [P3]. All these infinite family of manifolds were constructed from the elliptic surfaces $E(1)$ and $E(2)$ by applying the sequence of knot surgery in double nodes, ordinary blowups and rational blowdown. In this article, we obtain similar result starting from $E(n)$ for $n \geq 3$. Applying the sequence of knot surgery, ordinary blowups and rational blowdown we construct an infinite family of simply connected, nonsymplectic and pairwise nondiffeomorphic manifolds with nontrivial Seiberg-Witten invariants. Our motivation for constructing such manifolds comes from the following question raised by Ronald Stern.

Question. Which homotopy types of a simply connected, smooth, closed 4-manifold can be obtained from $E(n)$ by the sequence of knot surgery, ordinary blowups and rational blowdowns?

The author was informed by Jongil Park that a similar result has been obtained also by him and Ki-Heon Yun.

Acknowledgments: I am grateful to Ron Stern for many helpful discussions. I also wish to thank the referee for the helpful remarks.

1. ELLIPTIC FIBRATIONS

1.1. Elliptic fibrations as words in mapping class group. Let $f: X^4 \rightarrow S^2$ be a genus 1 Lefschetz fibration. Then X^4 is T^2 bundle over S^2 away from finitely many critical values of f . The fibers over these critical values are called singular fibers of the Lefschetz fibration. Such fibrations are characterized by their monodromy, i.e. a factorization of the identity element in the mapping class group M_1 of the fiber T^2 as a product of right-handed Dehn twists.

It is well-known that the mapping class group M_1 of the torus is generated by t_a, t_b of M_1 subject to the relations

$$t_a t_b t_a = t_b t_a t_b \quad \text{and} \quad (t_a t_b)^6 = 1$$

where t_a, t_b are Dehn twists along the standard curves a and b that generate the first homology of the torus.

Classification of genus 1 Lefschetz fibrations were given by Moishezon [M], who showed that after a possible perturbation such a fibration over S^2 is equivalent to one of the fibrations given by the words $(t_a t_b)^{6n} = 1$ in M_1 . The total space X of this fibration is $E(n)$, where $E(n)$ is a simply connected elliptic surface. It is known that $E(n)$ has holomorphic Euler characteristic $\chi_h = n$ with a section of $E(n) \rightarrow S^2$ that is a sphere with self-intersection $-n$.

1.2. Singular fibers in elliptic fibrations. In this section we give a brief introduction to certain singular fibers that occur in elliptic fibrations. We also prove the lemma that will be used in our construction. The complete treatment of the topology of elliptic surfaces and their singular fibers can be found in [HKK] and [KM].

Type I_1 : The fiber is an immersed 2-sphere with one positive double point. It has monodromy that is a conjugate of t_a and referred to as a fishtail fiber.

Type I_k : The singularity which has a monodromy conjugate of t_a^k ($k \geq 2$) is called an I_k singularity (or necklace fiber). Singular fibers of type I_k are a plumbing of k smooth 2-spheres of self-intersection -2 along a circle. I_2 singularity is also called a double node.

Lemma 1.1. *There exists an elliptic Lefschetz fibration on the surface $E(n)$ with a section, a singular fiber F of type I_{8n} , $(2n - 1)$ singular fibers $F_1, F_2, F_3, \dots, F_{2n-1}$ of type I_2 and two additional fishtail fibers.*

Proof. The word $(t_a t_b)^{6n} = 1$ in the mapping class group M_1 determines a genus 1 Lefschetz fibration on $E(n)$. Notice that by using the braid relation $t_a t_b t_a = t_b t_a t_b$, this word becomes equivalent to the following words

$$\begin{aligned} 1 &= (t_a t_b)^{6n} = t_a (t_b t_a t_b) (t_a t_b t_a) \dots (t_b t_a t_b) t_a t_b = \\ & t_a (t_a t_b t_a) (t_b t_a t_b) \dots (t_a t_b t_a) t_a t_b = t_a^2 (t_b t_a)^{6n-3} t_b t_a^2 t_b = \\ & t_a^2 (t_b t_a t_b) (t_a t_b t_a) \dots (t_a t_b t_a) t_b t_a^2 t_b = t_a^2 (t_a t_b t_a) (t_b t_a t_b) \dots (t_b t_a t_b) t_b t_a^2 t_b = \\ & t_a^3 (t_b t_a)^{6n-4} t_b^2 t_a^2 t_b = t_a^3 (t_b t_a t_b) (t_a t_b t_a) \dots (t_b t_a t_b) t_a t_b^2 t_a^2 t_b = \\ & t_a^3 (t_a t_b t_a) (t_b t_a t_b) \dots (t_a t_b t_a) t_a t_b^2 t_a^2 t_b = t_a^4 (t_b t_a)^{6n-6} t_b t_a^2 t_b^2 t_a^2 t_b \end{aligned}$$

Applying the braid relation repeatedly many times as above, the word $(t_a t_b)^{6n} = 1$ becomes equivalent to

$$t_a^{4n} t_b t_a^2 t_b^2 t_a^2 t_b^2 t_a^2 t_b^2 \dots t_a^2 t_b = 1$$

Next, conjugating by powers of t_a in the last relation, we have the following word

$$t_a^{4n} t_b t_a^2 t_b^2 t_a^{-2} t_a^4 t_b^2 t_a^{-4} \dots t_a^{4n-2} t_b^2 t_a^{-(4n-2)} t_a^{4n} t_b t_a^{-4n} t_a^{4n} = 1$$

Finally, multiplying by t_a^{4n} from the left and t_a^{-4n} from the right in the last relation, we obtain the following word

$$t_a^{8n} t_b t_a^2 t_b^2 t_a^{-2} t_a^4 t_b^2 t_a^{-4} \dots t_a^{4n-2} t_b^2 t_a^{-(4n-2)} t_a^{4n} t_b t_a^{-4n} = 1$$

This word translates into the singular fiber of type I_{8n} , $(2n-1)$ I_2 fibers and two fisthtail fibers. $\square$

2. SEIBERG-WITTEN INVARIANTS

In this section we review the basics of Seiberg-Witten invariants introduced by Seiberg and Witten. In order to state theorems in the following sections, we will view the Seiberg-Witten invariant of a smooth 4-manifold as a multivariable Laurent polynomial [W]. Let us recall that the Seiberg-Witten invariant of a smooth closed oriented 4-manifold X with $b_2^+(X) > 1$ is an integer valued function which is defined on the set of $spin^c$ structures over X [W]. For simplicity we assume that $H_1(X, \mathbf{Z})$ has no 2-torsion. Then there is a one-to-one correspondence between the set of $spin^c$ structures over X and the set characteristic elements of $H^2(X, \mathbf{Z})$ as following: To each $spin^c$ structure $\mathfrak{s}$ over X corresponds a bundle of positive spinors $W_{\mathfrak{s}}^+$ over X . Let $c(\mathfrak{s}) \in H_2(X)$ denote the Poincaré dual of $c_1(W_{\mathfrak{s}}^+)$. Each $c(\mathfrak{s})$ is a characteristic element of $H_2(X; \mathbf{Z})$ (i.e. its Poincaré dual $\hat{c}(\mathfrak{s}) = c_1(W_{\mathfrak{s}}^+)$ reduces mod 2 to $w_2(X)$).

In this set up we can view the Seiberg-Witten invariant as integer valued function

$$\text{SW}_X : \{k \in H^2(X, \mathbf{Z}) | k \equiv w_2(TX) \pmod{2}\} \rightarrow \mathbf{Z}.$$

The Seiberg-Witten invariant SW_X is a diffeomorphism invariant and its sign depends on an orientation of

$$H^0(X, \mathbf{R}) \otimes \det H_+^2(X, \mathbf{R}) \otimes \det H^1(X, \mathbf{R}).$$

If $\text{SW}_X(\beta) \neq 0$, then we call β a *basic class* of X . It is a fundamental fact that the set of basic classes is finite. It can be shown that, if β is a basic class, then so is $-\beta$ with

$$\text{SW}_X(-\beta) = (-1)^{(e+\text{sign})(X)/4} \text{SW}_X(\beta)$$

where $e(X)$ is the Euler number and $\text{sign}(X)$ is the signature of X .

Let $\{\pm\beta_1, \dots, \pm\beta_n\}$ be the set of nonzero basic classes for X . Consider variables $t_\beta = \exp(\beta)$ for each $\beta \in H^2(X; \mathbf{Z})$ which satisfy the relations $t_{\alpha+\beta} = t_\alpha t_\beta$. We may then view the Seiberg-Witten invariant of X as the symmetric Laurent polynomial

$$\text{SW}_X = b_0 + \sum_{j=1}^n b_j (t_{\beta_j} + (-1)^{(e+\text{sign})(X)/4} t_{\beta_j}^{-1})$$

where $b_0 = \text{SW}_X(0)$ and $b_j = \text{SW}_X(\beta_j)$.

Example 1. Let $E(n)$ be a simply connected minimally elliptic surface with holomorphic Euler characteristic $\chi = n$ and with no multiple fibers. Then we have $\text{SW}_{E(n)} = (t - t^{-1})^{n-2}$ where $t = \exp(T)$ and T is the cohomology class Poincare dual to the fiber class. Thus $\text{SW}_{E(n)}((n-2i)T) = (-1)^{i-1} \binom{n-2}{i-1}$ for $i = 1, \dots, n-1$ and $\text{SW}_{E(n)}(\beta) = 0$ for any other β .

Theorem 2.1. (Taubes) [T] Suppose that X is a closed symplectic 4-manifold with $b_2^+(X) > 1$. If K_X is a canonical classes of X , then $\text{SW}_X(\pm K_X) = \pm 1$.

3. KNOT SURGERY

Let X be a 4-manifold (with $b_2^+(X) > 1$) which contains a homologically-nontrivial torus T of self-intersection 0. Let $N(K)$ be a tubular neighborhood of K in S^3 , and let $T \times D^2$ be a tubular neighborhood of T in X . Then the knot surgery manifold X_K is defined by

$$X_K = (X \setminus (T \times D^2)) \cup (S^1 \times (S^3 \setminus N(K)))$$

Fintushel and Stern proved the theorem that shows Seiberg-Witten invariants of X_K can be completely determined by the Seiberg-Witten invariant of X and the Alexander polynomial of K [FS2]. Furthermore, if X and $X \setminus T$ are simply connected, then so is X_K .

Theorem 3.1. [FS2] *Assume that T lies in a cusp neighborhood in X , then Seiberg-Witten invariants of X_K is $\mathcal{SW}_{X_K} = \mathcal{SW}_X \cdot \Delta_K(t)$. If the Alexander polynomial $\Delta_K(t)$ of knot K is not monic then X_K admits no symplectic structure.*

We refer the reader to [FS3] for recent technique of Fintushel-Stern on knot surgery in double nodes.

4. RATIONAL BLOW-DOWNS AND ITS GENERALIZATIONS

In early nineties, the rational blow-down surgery was introduced by Fintushel and Stern [FS1]. The basic idea is that if a smooth 4-manifold X contains a particular configuration C_p of transversally intersecting 2-spheres whose boundary is the lens space $L(p^2, 1-p)$, then one can replace C_p with rational ball B_p to construct a new manifold X_p . If one knows the Donaldson and the Seiberg-Witten invariants of the original manifold X , then Fintushel-Stern show how one can determine same invariants of X_p . Initially Fintushel and Stern used this surgery technique to compute gauge invariants for simply connected regular elliptic surfaces with multiple fibers, i.e. $E(n)_{p,q}$ where p and q are relatively prime. Later this method has proven useful in the construction of many other interesting 4-manifolds. This rational blow-down surgery was generalized by Jongil Park [P1]. Below we discuss his generalized rational blow-down. Let $p \geq q \geq 1$ and p, q are relatively prime. Let $C_{p,q}$ be the smooth 4-manifold obtained by plumbing disk bundles over the 2-sphere according to the following linear diagram

$$\begin{array}{ccccccc} -r_k & -r_{k-1} & & \dots & & -r_1 \\ \bullet & \bullet & & & & \bullet \\ u_k & u_{k-1} & & & & u_1 \end{array}$$

where $p^2/(pq-1) = [r_k, r_{k-1}, \dots, r_1]$ is the unique continued linear fraction with all $r_i \geq 2$ and each vertex u_i of the linear diagram represents a disk bundle over 2-sphere with Euler number $-r_i$. According to Casson and Harer [CH], the boundary of $C_{p,q}$ is the lens space $L(p^2, 1-pq)$ which also bounds a rational ball $B_{p,q}$ with $\pi_1(B_{p,q}) = \mathbf{Z}_p$ and $\pi_1(\partial B_{p,q}) \rightarrow \pi_1(B_{p,q})$ surjective. If $C_{p,q}$ is embedded in a 4-manifold X then the generalized rational blowdown manifold $X_{p,q}$ is obtained by replacing $C_{p,q}$ with $B_{p,q}$, i.e., $X_{p,q} = (X \setminus C_{p,q}) \cup B_{p,q}$. If X and $X \setminus C_{p,q}$ are simply connected, then so is $X_{p,q}$. Notice that the case when $q = 1$ is the construction of Fintushel-Stern with $C_p = C_{p,1}$ given by

$$\begin{array}{ccccccc} -(p+2) & -2 & & \dots & & -2 \\ \bullet & \bullet & & & & \bullet \\ u_{p-1} & u_{p-2} & & & & u_1 \end{array}$$

Lemma 4.1. *Let $X_{p,q}$ be the manifold obtained from X by a rational blow-down of the configuration $C_{p,q}$. Then $b_2^+(X_{p,q}) = b_2^+(X)$ and $c_1^2(X_{p,q}) = c_1^2(X) + k$*

Proof. Notice that the manifold $C_{p,q}$ is negative definite, so that $b_2^+(X_{p,q}) = b_2^+(X)$. Using the fact that $c_1^2 = 3\sigma + 2e$, we have $c_1^2(X_{p,q}) = 3\sigma(X_{p,q}) + 2e(X_{p,q}) = 3(\sigma(X) + k) + 2(e(X) - k) = c_1^2(X) + k$

□

Theorem 4.2. [P1]. *Suppose X is a smooth 4-manifold with $b_2^+(X) > 1$ which contains a configuration $C_{p,q}$. If L is a characteristic line bundle on X such that, $SW_X(L) \neq 0$, $(L|_{C_{p,q}})^2 = -b_2(C_{p,q})$ and $c_1(L|_{L(p^2, 1-pq)}) = mp \in \mathbf{Z}_{p^2} \cong H^2(L(p^2, 1-pq); \mathbf{Z})$ with $m \equiv (p-1) \pmod{2}$, then L induces a SW basic class $\bar{L}$ of $X_{p,q}$ such that $SW_{X_{p,q}}(\bar{L}) = SW_X(L)$.*

Corollary. 3.2. [P3]. *Suppose X is a smooth 4-manifold with $b_2^+(X) > 1$ which contains a configuration $C_{p,q}$. If L is a SW basic class of X satisfying $L \cdot u_i = (r_i - 2)$ for any i with $1 \leq i \leq k$ (or $L \cdot u_i = -(r_i - 2)$), then L induces a SW basic class $\bar{L}$ of $X_{p,q}$ such that $SW_{X_{p,q}}(\bar{L}) = SW_X(L)$.*

5. CONSTRUCTION OF EXOTIC MANIFOLDS

In this section, we construct a family of simply connected, nonsymplectic and nondiffeomorphic manifolds starting from the elliptic surfaces $E(n)$ for $n \geq 3$ and applying the combination of knot surgery, blowup and rational blowdown. Using the Seiberg-Witten invariants, we distinguish their smooth structures.

Theorem 5.1. *There are infinitely many simply connected, nonsymplectic and pairwise nondiffeomorphic manifolds that can be obtained from elliptic surfaces $E(n)$ $n \geq 3$ by the sequence of knot surgery in double nodes, ordinary blow ups and rational blowdowns with (χ_h, c_1^2) given by following formulas.*

$$\left\{ \begin{array}{ll} \chi_h = 3, c_1^2 \leq 16 \\ \chi_h = 4, c_1^2 \leq 23 \\ \chi_h = 5, c_1^2 \leq 30 \\ \chi_h = 6, c_1^2 \leq 36 \\ \chi_h = n \geq 7, c_1^2 \leq 25k - 2 & \text{if } n = 4k \\ \chi_h = n \geq 7, c_1^2 \leq 25k + 5 & \text{if } n = 4k + 1 \\ \chi_h = n \geq 7, c_1^2 \leq 25k + 11 & \text{if } n = 4k + 2 \\ \chi_h = n \geq 7, c_1^2 \leq 25k + 18 & \text{if } n = 4k + 3 \end{array} \right.$$

Proof. Let first outline the general procedure how construct such family of simply connected, nonsymplectic 4-manifolds. According to lemma 1.1 $E(n)$

has elliptic fibration with $(2n+2)$ singular fibers, which are an I_{8n} singularity, $(2n-1)$ number of I_2 singularity and two fishtail fibers. First perform a knot surgery in s of the double nodes [FS2] of the $(2n-1)$ I_2 singularity using non fibered twist knots $K_1, K_2, \dots, K_s$, where $s = \lfloor \frac{7n+3}{4} \rfloor$. We denote the resulting manifold as $Y_{K_1, K_2, \dots, K_s}$. The manifold $Y_{K_1, K_2, \dots, K_s}$ is simply connected and has pseudo section S [FS2] which is sphere of self-intersection $-n$ with s positive double points. The pseudo section S intersects only one of the -2 spheres of I_{8n} singularity. To construct our manifolds, we blow up s double points on S to get sphere S' of self-intersection $-(n+4s)$ in $Y_{K_1, K_2, \dots, K_s} \# s \overline{\mathbf{CP}}^2$. Next blowup one of the fishtails if needed and smooth out its intersection with S' to get sphere S'' of the self-intersection $-(8n+3)$. Then remove one of the spheres from the I_{8n} configuration to get $(8n-1)$ linear chain of -2 spheres which together with S'' form the configuration C_{8n+1} . By blowing down of this configuration we get our manifolds. For simplicity let assume $K_1 = K_2 = K_3 = \dots = K_r = T(r)$ where $T(r)$ is non fibered r -twist knot. We will denote the manifold $Y_{K_1, K_2, \dots, K_s}$ in this set up as $Y(n)_{r,s}$ and the manifold obtained after rational blowdown as $Z(n)_{s,r,m}$, where m denotes the number of the blowups applied. Notice that above construction is well defined once $n \geq 7$. The case when $n < 7$ can be treated similarly with small modification of the construction above and by rationally blowing down the configurations of smaller length.

Let first assume that $n \geq 7$.

Case 1. $n = 4k$. By Lemma 1.1, there is fibration with I_{32k} singularity and $(2n-1) = 8k-1$ number of I_2 singularity and two fishtail fibers. Perform the knot surgery in $7k$ double nodes, blow up $s = 7k$ double points to get a sphere S' with self-intersection $-32k$. Next blowup one of the fishtails, smooth out its intersection with S' to get sphere S'' with self-intersection $-(32k+2)$. Use the linear chain of $(32k-2)$ of -2 spheres from I_{32k} singularity together with S'' to get the configuration of C_{32k} inside of $Y(n)_{r,s} \# (7k+1) \overline{\mathbf{CP}}^2$. By rationally blowing down this configuration we get manifolds with $c_1^2 = 25k - 2$ $\chi_h = 4k$.

Case 2. $n = 4k + 1$ By Lemma 1.1, there is fibration of $E(n)$ with one I_{32k+8} singularity and $(2n-1) = 8k+1$ number of I_2 singularity and two fishtail fibers. After performing the knot surgery in $s = 7k+2$ double nodes, blow up the double points of pseudo-section, one fishtail fiber and smooth out intersection of resulting sphere of self-intersection $-(32k+9)$ with fishtail fiber one gets the configuration of C_{32k+9} inside of $Y(n)_{r,s} \# (7k+3) \overline{\mathbf{CP}}^2$. After rationally blowing down this configuration, we get manifolds with $c_1^2 = 25k + 5$ and $\chi_h = 4k + 1$.

Case 3. $n = 4k + 2$. By Lemma 1.1, there is fibration with I_{32k+16} singularity and $(2n-1) = 8k+3$ number of I_2 singularity and two fishtail fibers. Perform knot surgery in $7k+4$ double nodes, blow up $s = 7k+4$ times to get a sphere S' with self-intersection $-(32k+18)$. Next use the linear

chain of $(32k + 14)$ of -2 spheres from I_{32k+16} singularity together with S' to get configuration of C_{32k+16} inside of $Y(n)_{r,s} \# (7k + 4) \overline{\mathbf{CP}}^2$. Rational blowdown of this configuration gives the manifolds with $c_1^2 = 25k + 11$ $\chi_h = 4k + 2$

Case 4. $n = 4k + 3$. Again by Lemma 1.1, $E(n)$ has fibration with I_{32k+24} singularity and $(2n - 1) = 8k + 5$ number of I_2 singularity and two fishtails. We perform knot surgery on $s = 7k + 6$ double nodes and blow up double points on pseudo-section to get configuration of $C_{32k+25} = C_{8n+1}$ configuration inside of $Y(n)_{r,s} \# (7k + 6) \overline{\mathbf{CP}}^2$. By rational blowdown we get manifold with $c_1^2 = 25k + 18$ and $\chi_h = 4k + 3$

Notice that these manifolds are simply-connected due to the fact that there is a fishtail fiber that was not used in the construction. By computing the Seiberg-Witten invariants, we will distinguish the different smooth structures constructed.

To compute Seiberg-Witten invariants of the manifold $Y(n)_{r,s}$, one applies the result of Fintushel-Stern on how Seiberg-Witten invariants change under the knot surgery [FS2]. Using the facts that the Seiberg-Witten function of $E(n)$ is $(e^T - e^{-T})^{n-2}$ and the Alexander polynomial of the twist knot $T(r)$ is $\Delta_{T(r)} = (2r - 1) + rt - rt^{-1}$, one computes the Seiberg-Witten invariants of $SW_{Y_{n,r}} = (e^T - e^{-T})^{n-2}((2r - 1) + re^{2T} - re^{-2T})^s$. By blow up formula for the Seiberg-Witten function [FS2], we have $SW_{Y(n)_{r,s} \# m \overline{\mathbf{CP}}^2} = SW_{Y_{n,r}} \cdot \prod_{j=1}^m (e^{E_j} + e^{-E_j})$, where E_j is an exceptional class coming from the j^{th} blow up. Furthermore, using Corollary 3.2 and Theorem 1.2 [P1], we compute the Seiberg-Witten invariants of $Z(n)_{s,r,m}$: Up to sign it has only one basic class which corresponds to top classes $\pm((n + 2s - 2)T + E_1 + \dots + E_m)$ and the value of the Seiberg-Witten function on these classes are $\pm r^s$. Now by applying the theorem of Taubes from Section 2, we detect that these manifolds are not symplectic. Since the Seiberg-Witten invariants are diffeomorphism invariants, the manifolds constructed yield an infinite family of simply connected, nonsymplectic and pairwise nondiffeomorphic manifolds once we consider different non fibered twist knots in the construction.

Case $n < 7$. We carry out the computation for the cases $n = 3, 4$. The remaining cases are similar.

i. $n = 3$. According to Lemma 1.1, $E(3)$ have one I_{24} singularity, 5 I_2 singularity and two fishtail fibers. Perform knot surgery in all 5 double nodes, blowup 5 double points of the pseudo section S , one of the fishtails and smooth out its intersection with the pseudo section to get sphere S' of self intersection -25 . By using 21 of the -2 spheres of the I_{24} singularity and S' one gets the configuration of C_{23} inside of $E(3) \# 6 \overline{\mathbf{CP}}^2$. After rational blowdown of C_{23} , one gets simply-connected manifolds with $c_1^2 = 16$, $\chi_h = 3$.

ii. $n = 4$. Again, by Lemma 1.1, $E(4)$ admits a fibration with one I_{32} singularity, 7 I_2 singularity and two fishtail fibers. Perform knot surgery in all 7 double nodes, blowup 7 double points of the pseudo section S , one of the fishtails and smooth out its intersection with the pseudo section to get sphere S' of self intersection -34 . By using 30 of the -2 spheres of the I_{32} singularity and S' one gets the configuration of C_{32} inside of $E(4)\#8\overline{\mathbf{CP}}^2$. After rational blowdown of C_{32} , we get simply-connected manifolds with $c_1^2 = 23$, $\chi_h = 4$.

The computation of Seiberg-Witten invariants is similar.

Now varying the number of double nodes s from 0 to $[7n + 3/4]$ and applying the blow ups, we can fill all the lattice points given as in the statement of the theorem. Notice that the number of basic classes will increase and c_1^2 reduce by blow up operation.

□

REFERENCES

- [CH] A. Casson and J. Harer, *Some homology lens spaces which bound homology rational ball*, Pacific.J. Math. **96** (1981), 23–36
- [GS] R. Gompf and A. Stipsicz, *4-manifolds and Kirby calculus*, Grad. Stud. in Math. AMS (1999).
- [FS1] R. Fintushel and R. Stern, *Rational blowdowns of smooth 4-manifolds*, Jour. Diff. Geom. **46** (1997), 181–235.
- [FS2] R. Fintushel and R. Stern, *Knots, links, and 4-manifolds*, Invent. Math. **134** (1998), 363–400.
- [FS3] R. Fintushel and R. Stern, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* , to appear in J. Amer. Math. Soc. **19** (2006), 171–180.
- [HKK] J. Harer, A. Kas and R. Kirby, *Handlebody decompositions of complex surfaces*, Mem.Amer.Math.Soc. **62** (1986), no.350.
- [KM] R. Kirby and P. Melvin, *The E_8 manifolds, singular fibers and handlebody decomposition*, Algebr. Geom. Topology **3** (2003), 577–568.
- [M] B. Moishezon, *Complex surfaces and connected sums of complex projective planes*, Lecture Notes in Mathematics **603** Springer-Verlag (1977)
- [P1] J. Park, *Seiberg-Witten invariants of generalized rational blow-downs*, Bull. Austral. Math. Soc. **56** (1997) 363–384
- [P2] J. Park, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math., **159** (2005), 657–667.
- [P3] J. Park, *Exotic smooth structures on $3\mathbf{CP}^2\#8\overline{\mathbf{CP}}^2$* , preprint.
- [PSS] J. Park, A. Stipsicz and Z. Szabo, *Exotic smooth structures on $\mathbf{CP}^2\#5\overline{\mathbf{CP}}^2$* , Math. Res. Letters **12** (2005), 701 – 712.
- [SS1] A. Stipsicz and Z. Szabo, *An exotic smooth structure on $\mathbf{CP}^2\#6\overline{\mathbf{CP}}^2$* , Geometry and Topology **9** (2005), 813–832.
- [SS2] A. Stipsicz and Z. Szabo, *Small exotic 4 manifolds with $b_2^+ = 3$* , preprint, <http://front.math.ucdavis.edu/math.GT/0411258>
- [T] C. Taubes, *The Seiberg-Witten invariants and symplectic forms*, Math. Res. Letters **1** (1994), 809–822.
- [W] E. Witten, *Monopoles and four-manifolds*, Math. Res. Letters **1** (1994), 769–796.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT IRVINE, IRVINE, CA
92697-3875

E-mail address: `aakhmedo@math.uci.edu`